

Detecting X-ray QPOs in Active Galaxies

Simon Vaughan^a and Philip Uttley^b

^a*X-Ray and Observational Astronomy Group, University of Leicester, Leicester,
LE1 7RH, UK*

^b*Laboratory for High Energy Astrophysics, Code 662, NASA Goddard Space Flight
Center, Greenbelt Road, Greenbelt, MD 20771, USA*

Abstract

Since *EXOSAT* produced the first good quality X-ray light curves of Seyfert galaxies there have been several claims of quasi-periodic oscillations (QPOs). None of these have withstood repeated analyses and observations. We review some problems concerning the detection of QPOs.

Key words: Active galactic nuclei, Time Series Analysis, periodic oscillations

1 Introduction: Why look for QPOs in Active Galaxies?

Active Galactic Nuclei (AGN) show strong, erratic brightness fluctuations in the X-ray band (Lawrence et al. 1987). At least for the Seyfert galaxies, the power spectrum of these variations is a power law-like continuum with a steep slope, except in a few cases where the slope can be seen to flatten at low frequencies (see e.g. Uttley et al. 2002; Markowitz et al. 2003). Processes with steep power spectra like this are known as *red noise* (Press 1978).

The X-ray variations in Seyfert galaxies bear more than a passing resemblance to those of X-ray binaries (XRBs; e.g. van der Klis 1995). In addition to a relatively featureless red noise continuum, XRBs often show QPOs in the form of peaks above the continuum (red noise) power spectrum. By analogy with XRBs, AGN may also possess QPOs. For example, assuming a $1/M_{BH}$ scaling of frequencies we might expect to see analogues of the high frequency QPOs of XRBs at $\sim 3 \times 10^{-3} (M_{BH}/10^6 M_{\odot})^{-1}$ Hz (Abramowicz et al., 2004). The detection of periodic or quasi-periodic variations from a Seyfert galaxy would

Email address: sav2@star.le.ac.uk (Simon Vaughan).

be a major discovery and could lead to a breakthrough in our understanding if the characteristic (peak) frequency could be identified with some physically meaningful frequency.

2 Textbook search methods

One of the most popular tools for time series analysis in astronomy is the *periodogram* (Schuster, 1898) which is simply the modulus-squared of the discrete Fourier transform of the time series (with an appropriate normalisation). This measures the amount of variability *power* as a function of temporal frequency. A concentration of power in a narrow range of frequencies would indicate a QPO¹. The notorious problem with the periodogram of noise processes is that the ordinates are distributed like χ^2 (with two degrees of freedom) about the true spectrum (Priestley, 1981). This means there are large fluctuations in the periodogram that can easily be mistaken for genuine peaks (periodic signals). Tests for periodic variations against a background of white (flat spectrum) noise that take account of these intrinsic sampling fluctuations are well established: see Fisher (1929) or section 6.1.4 of Priestley (1981), and also Leahy et al. (1983), van der Klis (1989) and Press et al. (1996). If the underlying source variability is not white noise the problem is more difficult and these methods will yield unreliable results (van der Klis 1989; Benlloch et al. 2001).

3 Possible sources of error

3.1 Systematic/instrumental effects

With non-imaging detectors (e.g. *EXOSAT*, *Ginga*, *RXTE*) there is always the possibility the light curve of a particular AGN is contaminated by other bright X-ray sources within the field of view. This possibility was realised in the case of NGC 6814 which had been identified as the best candidate for a Seyfert galaxy with an X-ray period based on non-imaging *EXOSAT* observations (Mittaz & Branduardi-Raymont 1989). Subsequent X-ray imaging observations made with *ROSAT* showed the periodic oscillations belonged to another source (V1432 Aquilae, a Cataclysmic Variable) located 37 arcmin from NGC 6814 (Madejski et al. 1993; Staubert et al. 1994). It is also possible for systematic, periodic changes in the detector properties to artificially modulate

¹ In the following we shall denote any strictly or quasi-periodic oscillation with the generic label “QPO.”

the observed light curve. An example is the *EXOSAT* data of NGC 5548. Papadakis & Lawrence (1993) reported significant periodicities in this moderate luminosity Seyfert but a reanalysis of the same data by Tagliaferri et al. (1996) demonstrated that one observation was affected by detector problems. The likelihood of these problems is greatly reduced with the current generation of low-background, imaging X-ray missions.

3.2 *Inappropriate null hypothesis: Assuming white noise*

Since the days of *EXOSAT* we have known that the X-ray variations of Seyfert galaxies have an intrinsically red power spectrum. This means the periodogram is not flat in expectation (as for white noise) and is generally not known *a priori*. As a result it is meaningless to apply the standard ‘periodogram peak’ tests (Fisher 1929; Leahy et al. 1983; Press et al. 1996). These all assume a null hypothesis of independent (white), Gaussian noise. Thus a high rejection probability for the null hypothesis obtained from any of these tests merely rejects white noise as the sole origin of the variance. It does not reject the more likely possibility that there are intrinsic red noise variations in the light curve (i.e. non-white, but also not periodic variations). Even a slight amount of intrinsic red noise variance will raise the spectrum above the white noise expectation and lead to spurious detections.

For example, Iwasawa et al. (1998) discussed a possible periodicity in IRAS 18325 – 5926 from a long (~ 5 day) *ASCA* observation. The significance of the candidate periodicity was estimated by testing against the hypothesis of purely Poisson noise. However, once the red noise variations intrinsic to the source were included in the periodogram fitting the candidate periodicity was no longer significant at the 95% level (Vaughan 2005). Other Seyfert galaxies for which candidate QPOs were reported without rigorously testing an origin in red noise include RX J0437.4 – 4711 (Halpern & Marshall 1996), MCG–6-30-15 (Lee et al. 2000), Mrk 766 (Boller et al. 2001) and IRAS 13224 – 3809 (Pfefferkorn et al. 2003).

3.3 *Subtle power spectral effects*

There are a number of ways that one can account for red noise in a light curve. But this is often not a trivial task and there are many possibilities for overestimating the significance of candidate QPOs. For example, Fiore et al. (1989) reported candidate QPOs in NGC 4151 based on three *EXOSAT* observations. The authors claimed high ($> 99\%$) significance peaks in the periodograms after the data were “pre-whitened” by detrending the light curves using a parabola. However, no account was made of the uncertainties involved in this




Fig. 1. Periodogram of the 1986/060 *EXOSAT* observation of NGC 4151 measured from the background-subtracted 1 – 9 keV ME data binned to 120 s (obtained through the *EXOSAT* data archive at HEASARC/GSFC). The histogram shows the raw data and the solid curve marks the Maximum Likelihood fit using a model comprising a power law (red noise) plus constant (Poisson noise). The dashed line represents a (minimum) 95% confidence limit for the detection of a QPO calculated as a factor $\gamma_\epsilon = -\ln[1 - (1 - \epsilon)^{1/n}]$ above the model, where n is the number of frequencies and ϵ is the ‘false alarm probability’. This quick calculation slightly underestimates the threshold because it does not account for uncertainties in the modelling (see Vaughan 2005). The arrow indicates the position of the candidate QPO discussed by Fiore et al. (1989).

procedure, which would undoubtedly leave some residual red noise component in the spectrum. If the continuum power at low frequencies (where the QPOs occurred and where the red noise will be strongest) was underestimated by only a small fraction this would drastically inflate the significance of the candidate QPOs. Figure 1 shows a re-analysis of one of these datasets. Explicitly fitting the red noise and Poisson noise components of the spectrum allows us to place more realistic limits on the significances of any QPOs, which now fall below the 95% confidence detection threshold.

The above example is one where the uncertainty in modelling the aperiodic power spectrum (i.e. the intrinsic red noise and instrumental Poisson noise) led to an artificially enhanced periodicity significance. In general, not accounting for the uncertainty in the null hypothesis model (the continuum) often tends to overestimate the significance of any peaks in the residual (“whitened”) periodogram. For example, in their re-analysis of the *EXOSAT* observations of NGC 5548, Tagliaferri et al. (1996) found that in every case the significance of the candidate QPO was lower than previously reported ($\leq 95\%$) once due allowance was made for the uncertainties in modelling the spectrum.

It is also possible to overestimate the significance of a periodogram peak when the periodogram is oversampled. Halpern et al. (2003) identified possible QPOs in three Seyfert 1 galaxies (Ton S180, RX J0437.4 – 4711, 1H

0419-577) based on a uniform analysis of all the available long *EUVE* observations. The significances of the periodicities were estimated using a Monte Carlo procedure similar to that discussed by Benlloch et al. (2001). However, the periodograms were oversampled (computed at frequencies between, as well as at, the standard Fourier frequencies). This increases the number of frequencies searched and therefore also the chance of seeing a spurious signal (Vaughan 2005).

Finally there are cases where genuinely significant power spectral features have been detected but were not caused by periodic oscillations. Papadakis & Lawrence (1995) reported a broad bump in the power spectrum of NGC 4051 (based on *EXOSAT* observations). However, recent observations with *ROSAT*, *RXTE* and *XMM-Newton* (Green et al. 1999; McHardy et al. 2004) have provided much tighter constraints on the power spectrum and have shown that the underlying continuum spectrum of NGC 4051 breaks at a similar frequency to that reported by Papadakis & Lawrence (1995). The broad bump in the spectrum reported from the *EXOSAT* observation is therefore likely to be due to the break in the continuum spectrum rather than an additional QPO feature.

4 Conclusions

A major source of ‘false alarm’ QPOs is the use of an inappropriate continuum power spectrum. Arbitrarily assuming a white noise spectrum, or red noise with an *ad hoc* slope, will alter the apparent strength (significance) of any peaks in the residual periodogram. The continuum spectrum must be modelled explicitly. Furthermore, one must account for uncertainties in the continuum modelling. Failure to include these may result in an overly optimistic significance calculation (Israel & Stella 1996; Tagliaferri et al. 1996; Vaughan 2005).

The above arguments apply not only to searching for peaks in the periodogram but also for searches based on *epoch folding*, whereby one bins the time series into phase bins at a test period. At the correct period the periodic variations will sum while any background noise will cancel out, thus revealing the profile of the periodic pulsations. However, the background noise will only cancel out if it is temporally independent, i.e. white noise. Again, the presence of any underlying red noise variations will produce unreliable results unless checked with e.g. Monte Carlo simulations² (Benlloch et al. 2001).

² A further word of caution is required here. For the correct application of Monte Carlo methods it is crucial to use a random noise generator that correctly produces the statistical aspects of red noise and the observational sampling, otherwise the significances may again be badly biased (Timmer & König 1995).

References

- Abramowicz, M. A., Kluźniak, McClintock J. E., et al., 2004 “The Importance of Discovering a 3:2 Twin-Peak Quasi-periodic Oscillation in an Ultraluminous X-Ray Source, or How to Solve the Puzzle of Intermediate-Mass Black Holes” 2004, *ApJ*, 609, L63-L65
- Benlloch, S., Wilms J., Edelson R., et al., 2001, “Quasi-periodic Oscillation in Seyfert Galaxies: Significance Levels. The Case of Markarian 766” *ApJ*, 562, L121-L124
- Boller, Th., Keil R., Trümper J., et al., 2001, “Detection of an X-ray periodicity in the Narrow-line Seyfert 1 Galaxy Mrk 766 with XMM-Newton” *A&A*, 365, L146-L151
- Fiore F., Massaro E., Perola G. C., et al. 1989, “Small-amplitude coherent signals in the X-ray emission of NGC 4151” *ApJ*, 347, 171-178
- Fisher, R. A., 1929, “Tests of significance in harmonic analysis” *Proc. Royal Society of London: Series A*, 125, 54-59
- Green A. R., McHardy I. M., Done C., 1999, “The discovery of non-linear X-ray variability in NGC 4051” *MNRAS*, 305, 309-318
- Halpern J. P., Marshall H. L. 1996, “A Long EUVE Observation of the Seyfert Galaxy RX J0437.4-4711” *ApJ*, 464, 760-764
- Halpern J. P., Leighly K. M., Marshall H. L. 2003, “An Extreme Ultraviolet Explorer Atlas of Seyfert Galaxy Light Curves: Search for Periodicity” *ApJ*, 585, 665-676
- Israel, G. L., Stella, L., 1996, “A New Technique for the Detection of Periodic Signals in ‘Colored’ Power Spectra” *ApJ*, 468, 369-379
- Iwasawa, K., Fabian A. C., Brandt W. N., et al., 1998, “Detection of an X-ray periodicity in the Seyfert galaxy IRAS 18325-5926” *MNRAS*, 295, L20-L24
- Lawrence, A., Watson M. G., Pounds K. A., Elvis M., 1987, “Low-frequency divergent X-ray variability in the Seyfert galaxy NGC4051” *Nature*, 325, 694-696
- Leahy, D. A., Darbro W., Elsner R. F., et al., 1983, “On searches for pulsed emission with application to four globular cluster X-ray sources - NGC 1851, 6441, 6624, and 6712” *ApJ*, 266, 160-170
- Lee J. C., Fabian A. C., Reynolds C. S., et al., 2000, “The X-ray variability of the Seyfert 1 galaxy MCG-6-30-15 from long ASCA and RXTE observations” *MNRAS*, 318, 857-874
- Madejski G. M. Done C., Turner T. J., et al., 1993, “Solving the Mystery of the X-Ray Periodicity in the Seyfert Galaxy NGC6814” *Nature*, 365, 626-628
- Markowitz A., Edelson R., Vaughan S., et al. 2003, “X-Ray Fluctuation Power Spectral Densities of Seyfert 1 Galaxies” *ApJ*, 593, 96-114
- McHardy I. M., Papadakis I. E., Uttley P., et al., 2004, “Combined long and short time-scale X-ray variability of NGC 4051 with RXTE and XMM-Newton” *MNRAS*, 348, 783-801
- Mittaz J. P. D., Branduardi-Raymont G., 1989, “The flux and spectral variability of NGC 6814 as observed with EXOSAT” *MNRAS*, 238, 1029-1046

- Papadakis I. E., Lawrence A., 1993, “Quasi-periodic oscillations in X-ray emission from the Seyfert galaxy NGC5548” *Nature*, 361, 233-236
- Papadakis I. E., Lawrence A., 1995, “A detailed X-ray variability study of the Seyfert galaxy NGC 4051” *MNRAS*, 272, 161-183
- Pfefferkorn F., Boller Th., Burwitz V., Predehl P., 2003, “Chandra observation of the narrow-line Seyfert 1 galaxy IRAS 13224-3809” in proceedings of the *MAXI workshop on AGN variability*, ed. N. Kawai, H. Negoro, A. Yoshida, T. Mihara, Seiyo Press (Tokyo), p137-141 (astro-ph/0106203)
- Press, W. H. 1978, “Flicker noises in astronomy and elsewhere” *Comments on Astrophysics*, 7, 103-119
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1996, *Numerical Recipes*, Cambridge Univ. Press. (Cambridge)
- Priestley M. B. 1981, *Spectral Analysis and Time Series*, Academic Press (London)
- Schuster A. 1898, “An investigation of hidden periodicities with application to a supposed 26-day period of meteorological phenomena” *Terrestrial Magnetism*, 3, 13-41
- Staubert R., König M., Friedrich S., et al., 1994, “On the newly discovered 3.4 hour modulated X-ray source (RXJ1940.1-1025) near NGC 6814” *A&A*, 288, 513-518
- Tagliaferri G., Bao G., Israel G. L., et al., 1996, “The soft and medium-energy X-ray variability of NGC 5548: a reanalysis of EXOSAT observations” *ApJ*, 465, 181-190
- Timmer J., König M. 1995, “On generating power law noise” *A&A*, 300, 707-710
- Uttley P., McHardy I. M., Papadakis I., 2002, “Measuring the broad-band power spectra of active galactic nuclei with RXTE” *MNRAS*, 332, 231-250
- van der Klis M. 1989, “Fourier techniques in X-ray timing” in *Timing Neutron Stars*, ed. H. Ogelman, E. P. J. van den Heuvel, Kluwer (Dordrecht), NATO ASI Series C 262, p27-69
- van der Klis M. 1995, “Rapid aperiodic variability in X-ray binaries” in *X-ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs, E. P. J. van den Heuvel, Cambridge Univ. Press (Cambridge), p252-307
- Vaughan S., 2005, “A simple test for periodic signals in red noise” *A&A*, 431, 391-403